

Opposite sign correlations in fermion or boson gases

Alex D. Gottlieb^{1,*} and Thorsten Schumm²

¹*Wolfgang Pauli Institute, Nordbergstrasse 15, A-1090 Vienna, Austria*

²*Atominstytut der Österreichischen Universitäten,
TU-Wien, Stadionallee 2, A-1020 Vienna, Austria*

We investigate pair correlations in trapped fermion and boson gases as a means to probe the quantum states producing the density fluctuations. We point out that “opposite sign correlations” (meaning pair correlations that are positive for fermions and negative for bosons) unambiguously indicate that the quantum many-particle state cannot be “free.” In particular, a system of fermions that exhibits positive pair correlations cannot be described by any Slater determinant wavefunction. This insight may help one to interpret results of current experiments on ultracold atomic gases.

PACS numbers: 03.75.Be, 42.50.Lc, 42.30.-d, 31.15.-p

In 1956 Hanbury Brown and Twiss performed a landmark experiment where they observed photon bunching in light emitted by a chaotic source [1]. The correlations they detected “stimulated the birth of modern quantum optics” [2, 3]. Forty years later, in another pioneering experiment, temporal Hanbury Brown-Twiss (HBT) correlations were observed for massive bosonic particles in a cold atomic beam, followed by the observation of antibunching in electron and neutron beams [4–6]. Bunching (or antibunching) may now be observed directly in position space when a boson (or fermion) gas is released from a trap [2, 3].

The last twelve years have witnessed rapid advances in the field of ultracold atoms, and there has naturally been great interest in the density correlations that can be measured in experiments on quantum gases of these atoms [7, 8]. Nowadays (2004-2007) correlations are being used to investigate ever more complex quantum states, e.g., squeezed number states of systems of indistinguishable atoms going through the Mott insulator transition in optical lattices [9, 10]. Momentum correlations are studied in collisions of two Bose-Einstein condensates and in the dissociation of weakly bound dimers [11, 12]. In weakly interacting low dimensional systems, where no clear phase transition to a condensed state takes place at low temperatures, correlations are used to characterize the degeneracy of the system [13]. In the strongly interacting regime, second- and third-order correlations at zero distance have been used to identify the Tonks-Girardeau gas [14, 15].

One may analyze pair correlations in order to

learn about the quantum states of trapped boson gases or atoms in optical lattices [7, 8]. The power of this kind of intensity interferometry as a tool to probe complex many-body quantum states of ultracold atoms is highlighted in a short article published in 2004, which considers models of superfluid fermion gases and optical lattice Mott insulators [16]. These models feature what we are calling opposite sign correlations, i.e., negative pair correlations for bosons, whose symmetry index is conventionally positive, and positive pair correlations for fermions, whose symmetry index is conventionally negative.

Our purpose here is to point out that if a many-boson or many-fermion system exhibits opposite sign correlations, then the state in question necessarily has a certain complexity. For example, consider a fermion gas of — say — 10^5 atoms. If the gas exhibits any positive pair correlations when it has been prepared in a certain state, then that state cannot be represented by a simple Slater determinant wavefunction. In general, if one probes a many-boson or many-fermion state and finds that it exhibits opposite sign correlations, then, even without any model for the unknown state, one may infer that it is not a “free” state, i.e., it does not have the form of a grand canonical ensemble for noninteracting indistinguishable particles. We believe that opposite sign correlations can be observed in current experimental setups and may even have already been observed and passed unnoticed [17].

Let us discriminate between correlations that are merely due to Bose or Fermi statistics, and

correlations that reflect a further complexity of the quantum state, the kind of complexity that demands for itself a *post*-Hartree or *post*-Hartree-Fock description. The bunching or antibunching observed in HBT effects [2–5, 9, 10] can be attributed to the symmetric or antisymmetric character of the many-particle wavefunctions that describe the system. For example, fermion antibunching can be explained if one considers the simplest many-fermion wavefunctions, i.e., Slater determinants, which are just antisymmetrized products of one-particle wavefunctions. Slater determinants necessarily feature the negative correlations that one sees in the (fermionic) HBT effect. However, more complex many-fermion wavefunctions, which must be represented by *superpositions* of two or more Slater determinants, can exhibit *positive* correlations. For example, let $|1\rangle, |2\rangle, |3\rangle$, and $|4\rangle$ represent four orthonormal one-fermion states and let $|1\rangle \wedge |2\rangle$ represent the two-particle Slater determinant formed from states $|1\rangle$ and $|2\rangle$. The two-particle state

$$\psi = \frac{1}{\sqrt{2}} |1\rangle \wedge |2\rangle + \frac{1}{\sqrt{2}} |3\rangle \wedge |4\rangle$$

does involve positive correlations of occupation number: occupation of state $|1\rangle$ is perfectly correlated with occupation of state $|2\rangle$. This state features correlations that cannot be explained by mere antisymmetry because ψ cannot be written as a single Slater determinant. To distinguish these kinds of correlations from simple (HBT) correlations, we refer to Slater determinant wavefunctions as “free” states and we call complex states like ψ “nonfree” [18, 19]. When quantum chemists and condensed matter physicists speak of “correlated” systems of electrons, they usually mean “nonfree” systems.

A free state is a state of a fermion or boson field in which occupation numbers, relative to some system of orthogonal one-particle modes, are statistically independent. Slater determinant wavefunctions represent free states since the one-particle orbitals that participate in the determinant are certainly occupied, while any orbital orthogonal to these is certainly unoccupied. The other fermionic free states are best represented by density matrices on the Fock space. All grand canonical thermal equilibrium ensembles of noninteracting fermions at positive temperature are examples of the latter. All number states in the fermion Fock space

are free states, but the only *pure* bosonic free state is the vacuum; all others are *mixed* states, which have to be represented by density matrices of rank greater than 1. Bosonic free states are called “gaussian” in the context of photon statistics, and are used to describe chaotic thermal radiation fields in quantum optics [20].

A free state is completely determined by its one-particle correlation operator (i.e., all first-order correlations, including off-diagonal correlations). Free states are conserved by free dynamics (where the particles do not interact with one another) and by efficient measurement of a one-particle observable. Therefore, if an ultracold gas is prepared in a free state and subjected to a time-dependent external potential, then, as long as particle-particle interactions may be disregarded, the state of the gas will remain free. In particular, switching off a trapping potential and allowing a gas to undergo “time-of-flight” should not destroy the freeness of its state.

The next few paragraphs only discuss systems of fermions; we’ll come back to bosons later.

If a many-fermion system is in a free state, and one then observes the positions of all the particles, the random pattern or “configuration” of points that results is a sample of a “determinantal point process” [21]. Determinantal point processes crop up unexpectedly in diverse areas of mathematics and physics, e.g., random matrix theory [22, 23] and graph theory [24], and they have been studied extensively [22, 24–28]. Although “it remains to investigate the processes thoroughly in connection to statistical inference” [29], determinantal point processes have statistical properties that can provide tests of “determinantality” and, hence, of “freeness.” Specifically, determinantal point processes have the property that density fluctuations in disjoint regions are never positively correlated, and therefore, when an experiment on ultracold fermions reveals positive density-density correlations, this can only mean that the state of the many-fermion system is not free.

A *point process* on a bounded region of space S is a random finite subset X of S . A point process X is *determinantal on S with kernel $\mathcal{K} : S \times S \rightarrow \mathbb{C}$* (which is assumed to be self-adjoint) if

$$\mathbb{E} \left[\prod_{j=1}^m \#(X \cap E_j) \right] \quad (1)$$

— the statistical expectation of the product of the occupation counts for arbitrary disjoint measurable subsets E_1, \dots, E_m of S — equals

$$\int_{E_1} \cdots \int_{E_m} \det (\mathcal{K}(x_i, x_j))_{i,j=1}^m dx_1 \cdots dx_m \quad (2)$$

for all $m \geq 1$ [27, 28]. From (1) = (2) it follows that, for a determinantal point process X , the numbers of points in any two disjoint regions of space are never positively correlated. That is, if R_1 and R_2 are two disjoint regions of space, and if $N_1 = \#(X \cap R_1)$ and $N_2 = \#(X \cap R_2)$ denote, respectively, the random number of points in each of these regions, then $\langle N_1 N_2 \rangle \leq \langle N_1 \rangle \langle N_2 \rangle$. Furthermore, in case the particle numbers in R_1 and R_2 are *uncorrelated*, i.e., when $\langle N_1 N_2 \rangle = \langle N_1 \rangle \langle N_2 \rangle$, the two restricted point processes $X \cap R_1$ and $X \cap R_2$ are *independent*. If one views a determinantal process through an “observation window” R [29], the restricted point process $X \cap R$ is also determinantal.

The configuration statistics of a free fermion state ω are determinantal with kernel

$$\mathcal{K}(x, y) = \omega(\hat{\psi}^\dagger(y)\hat{\psi}(x)),$$

where $\hat{\psi}(x)$ denotes the field operator at x . The kernel of the point process viewed through observation window R is simply

$$\mathbf{1}_R(x)\mathcal{K}(x, y)\mathbf{1}_R(y).$$

Therefore, when one observes the configuration X of a system of fermions in a free state ω , the random occupation numbers

$$\begin{aligned} N_1 &= \#(X \cap R_1) \\ N_2 &= \#(X \cap R_2) \end{aligned}$$

satisfy

$$\langle N_1 N_2 \rangle_\omega \leq \langle N_1 \rangle_\omega \langle N_2 \rangle_\omega \quad (3)$$

if R_1 and R_2 do not overlap. More generally,

$$\left\langle \prod_{i=1}^k N_i \right\rangle_\omega \leq \prod_{i=1}^k \langle N_i \rangle_\omega$$

if the the regions R_1, R_2, \dots, R_k are disjoint. Violations of the “antibunching” inequality (3) indicate that the many-fermion state ω is not free.

For bosons, it is *negative* pair correlations that indicate nonfreeness. A system of bosons cannot be in a free state if the density fluctuations at any pair of (distinct) points are negatively correlated. The point processes corresponding to the bosonic free states are known as “permanantal point processes” [25, 30] and these feature positive density-density correlations. That is, if X is the random configuration of bosons in a free state ω , and if $N_1 = \#(X \cap R_1)$ and $N_2 = \#(X \cap R_2)$, then

$$\langle N_1 N_2 \rangle_\omega \geq \langle N_1 \rangle_\omega \langle N_2 \rangle_\omega \quad (4)$$

whenever R_1 and R_2 disjoint, and therefore, violations of the “bunching” inequality (4) signify that the many-boson state ω is not free. On a quantum statistical level, the HBT correlations that may be observed in starlight are due to the thermal character of the source of that light, from which gaussian (i.e., free) photon fields are assumed to emanate [20, 31]. Incoherent thermal sources produce radiation fields whose pair correlations are positive or zero, but other kinds of light sources can produce photon fields with *negative* pair correlations [32].

Free states correspond to *grand canonical* ensembles of noninteracting bosons or fermions, but it is more appropriate to model thermal equilibrium states of ultracold gases by the *canonical* ensemble, for the trapped gas is a closed system of massive particles. However, when the number of particles is large, it is reasonable to suppose that canonical ensembles of free bosons (or fermions) exhibit *nearly* permanantal (or determinantal) statistics. Such “equivalence of ensembles” has recently been established rigorously for untrapped particles [33].

In conclusion, when opposite sign correlations are observed, this suggests that interactions between the particles, whether they be collisions or indirectly mediated interactions, are responsible for a certain structural complexity of the many-body state. Such opposite sign correlations should be detectable in absorption images of ultracold atomic gases. When opposite sign correlations are observed one may infer that the state in question is definitely not free. Remarkably, this inference does not require any knowledge or model of the unknown state.

Acknowledgements. *We thank A. Scrinzi and J. Schmiedmayer for very helpful instruction.*

A.G. is supported by the Austrian Ministry of Science via its grant for the Wolfgang Pauli Institute and by the Austrian Science Foundation (FWF) via the START Project Y-137-TEC.

-
- * Electronic address: www.alexgottlieb.com
- [1] R. Hanbury Brown and R. Q. Twiss, *Correlation between Photons in two Coherent Beams of Light*, Nature 177, 27 (1956)
 - [2] M. Schellekens, R. Hoppeler, A. Perrin, J. Viana Gomes, D. Boiron, A. Aspect, and C. I. Westbrook, *Hanbury Brown Twiss effect for ultracold quantum gases*, Science 310, 648-651 (2005)
 - [3] T. Jelte, J. M. McNamara, W. Hogervorst, W. Vassen, V. Krachmalnicoff, M. Schellekens, A. Perrin, H. Chang, D. Boiron, A. Aspect and C. I. Westbrook, *Hanbury Brown Twiss effect for bosons versus fermions*, Nature 445, 402-405 (2007)
 - [4] M. Yasuda and F. Shimizu, *Observation of Two-Atom Correlation of an Ultracold Neon Atomic Beam*, Phys. Rev. Lett. 77, 3090 - 3093 (1996)
 - [5] H. Kiessel, A. Renz and F. Hasselbach, *Observation of Hanbury Brown-Twiss anticorrelations for free electrons*, Nature 418, 392 - 394 (2002)
 - [6] M. Iannuzzi, A. Orecchini, F. Sacchetti, P. Facchi and S. Pascazio, *Direct Experimental Evidence of Free-Fermion Antibunching*, Phys. Rev. Lett. 96, 080402 (2006)
 - [7] M. Naraschewski and R. J. Glauber, *Spatial coherence and density correlations of trapped Bose gases*, Phys. Rev. A 59, 4595 - 4607 (1999)
 - [8] J. Grondalski, P. M. Alsing and H. Deutsch, *Spatial correlation diagnostics for atoms in optical lattices*, Optics Express 5, 249 - 261 (1999)
 - [9] S. Foelling, F. Gerbier, A. Widera, O. Mandel, T. Gericke and I. Bloch, *Spatial quantum noise interferometry in expanding ultracold atom clouds*, Nature 434, 481 - 484 (2005)
 - [10] T. Rom, Th. Best, D. van Oosten, U. Schneider, S. Flling, B. Paredes and I. Bloch, *Free fermion antibunching in a degenerate atomic Fermi gas released from an optical lattice*, Nature 444, 733 - 736 (2006)
 - [11] A. Perrin, H. Chang, V. Krachmalnicoff, M. Schellekens, D. Boiron, A. Aspect and C. I. Westbrook *Atom-atom correlations in spontaneous four wave mixing of two colliding Bose-Einstein Condensates*. arXiv:quant-ph/0704.3047 v1 (2007)
 - [12] M. Greiner, C. A. Regal, J.T. Stewart and D. S. Jin, *Probing pair-correlated fermionic atoms through correlations in atom shot noise*, Phys. Rev. Lett. 94, 110401 (2005)
 - [13] J. Esteve, J.-B. Trebbia, T. Schumm, A. Aspect, C. I. Westbrook, and I. Bouchoule, *Observations of Density Fluctuations in an Elongated Bose Gas: Ideal Gas and Quasicondensate Regimes*, Phys. Rev. Lett. 96, 130403 (2006)
 - [14] B. Laburthe Tolra, K. M. O'Hara, J. H. Huckans, W. D. Phillips, S. L. Rolston, and J. V. Porto, *Observation of Reduced Three-Body Recombination in a Correlated 1D Degenerate Bose Gas*, Phys. Rev. Lett. 92, 190401 (2004)
 - [15] T. Kinoshita, T. Wenger, and D. S. Weiss, *Local Pair Correlations in One-Dimensional Bose Gases*, Phys. Rev. Lett. 95, 190406 (2005)
 - [16] E. Altman, E. Demler and M. Lukin, *Probing many-body states of ultracold atoms via noise correlations*, Physical Review A 70, 013603 (2004)
 - [17] We have in mind Refs. 9 and 10 (see Figure 4d in both Letters).
 - [18] A. D. Gottlieb and N. J. Mauser, *New measure of electron correlation*, Phys. Rev. Lett. 95 (12), 123003 (2005)
 - [19] A. D. Gottlieb and N. J. Mauser, *Properties of nonfreeness: an entropy measure of electron correlation*, Submitted (2007)
 - [20] J. R. Klauder and E. C. G. Sudarshan. *Fundamentals of Quantum Optics*, W. A. Benjamin, Inc., New York, 1968
 - [21] A. D. Gottlieb, *Introduction to determinantal point processes from a quantum probability viewpoint*, To appear in Quantum Probability and Infinite Dimensional Analysis series, World Scientific.
 - [22] A. Soshnikov, *Determinantal random point fields*, Russian Math. Surveys 55, 923 - 975 (2000)
 - [23] K. Johansson, *Random matrices and determinantal processes*, Preprint arXiv:math.PR/0510038 v1 (2005)
 - [24] R. Lyons, *Determinantal probability measures*, Publ. Math. Inst. Hautes Études Sci. 98, 167 - 212 (2003)
 - [25] O. Macchi, *The coincidence approach to stochastic point processes*, Adv. Appl. Prob. 7, 83 - 122 (1975)
 - [26] T. Shirai and Y. Takahashi, *Random point fields associated with certain Fredholm determinants I: fermion, Poisson and boson point processes*, J. Functional Analysis 205, 414 - 463 (2003)
 - [27] J. Ben Hough, M. Krishnapur, Y. Peres and B. Virág, *Determinantal processes and independence*, Probability Surveys 3, 206 - 229 (2006)
 - [28] D. J. Daley and D. Vere-Jones, *An Introduction to the Theory of Point Processes*, Springer-Verlag, New York, 2003
 - [29] J. Møller and R. Waagepetersen, *Modern statistics for spatial point processes*, Preprint (2007) to appear in Scandinavian Journal of Statistics
 - [30] P. McCullagh and J. Møller, *The permanental*

- process*, Advances in Applied Probability 38, 873 - 888 (2006)
- [31] R. J. Glauber. *Photon correlations*, Phys. Rev. Lett. 10, 84 - 86 (1963)
- [32] See page 246 of Ref. 20, where this is explicit.
- [33] H. Tamura and K. R. Ito, *A canonical ensemble approach to the fermion/boson random point processes and its applications*, Communications in Mathematical Physics 263, 353 - 380 (2006)